Recent advances in peptic ulcer bleeding

Ian LP Beales

Address: Gastroenterology Department, Norfolk and Norwich University Hospital, Norwich, Norfolk NR4 7UZ, UK

Email: i.beales@uea.ac.uk

F1000 Medicine Reports 2009, 1:44 (doi:10.3410/M1-44)

The electronic version of this article is the complete one and can be found at: <http://F1000.com/Reports/Medicine/content/1/44>

Abstract

Peptic ulcer bleeding remains a common cause of hospital admission, morbidity and mortality. Data published since 2006 illustrate that assessment, endoscopic and pharmacological management, and follow-up strategies can be refined to improve the overall prognosis of peptic ulcer bleeding.

Introduction and context

Acute upper gastrointestinal bleeding remains an important medical emergency. A recent large nationwide audit in the UK (6,750 cases) showed that overall rates of admission to hospital were 103 out of 100,000 adults, with an overall mortality of 10% in new admissions and 26% in those who have a gastrointestinal haemorrhage whilst an inpatient. Peptic ulcer bleeding (PUB) remains the single most common cause of upper gastrointestinal haemorrhage (36% of cases) and the UK audit showed a mortality of 3.8% in patients under 60 years old rising to 10% in patients over 80 years old [1]. These results are rather worse than those reported in trials in specialist centres and probably reflect the increasing age and comorbidity of patients in the 'real world', as well as the relatively slow spread of optimal practice (for instance, only 6% of patients had dual endoscopic therapy [1]). Recent studies have further refined our management strategies for PUB and these can be viewed in terms of risk scoring and stratification, resuscitation, endoscopic therapy, pharmacotherapy and prevention.

Recent advances

Risk stratification and scoring

The Rockall score is widely used to compare outcomes and is very useful as an audit tool, having been validated in several populations [2]. It performs less well as a guide to management, especially in predicting very low-risk bleeders, who can safely be discharged without hospital admission. The Glasgow-Blatchford score (GBS) was

initially shown to be superior in defining this low risk group and a recent validation study in 676 separate patients confirmed that a score of 0 [pulse <100, systolic blood pressure >110 mmHg, urea <6.5 mmol/l, haemoglobin >13.0 g/dl (men) or 12.0 g/dl (women), and an absence of syncope, melaena, heart failure or liver disease] was associated with a zero rate of death, transfusion or endoscopic intervention [3]. A Canadian study utilised a modified GBS (omitting urea and syncope) and although effective, it did perform less well than the full score (5% re-bleeding rate in the 0 scoring group) [4]. Thus, early discharge strategies based on the full GBS scoring system should be implemented to increase efficient use of resources.

Resuscitation

Whilst prompt and appropriate fluid resuscitation is essential to the effective management of PUB, there are remarkably few data on which to base management and transfusion strategies and these remain essentially clinical decisions. An increasing number of patients with PUB have been treated with vitamin K antagonists such as warfarin, and the availability of prothrombin complex concentrate (which contains factors II, VII, IX and X) is a real advance in the reversal of anticoagulation in such patients [5]. Although never examined specifically in PUB, the rapid and effective reversal of anticoagulation achieved with this agent implies it should be central to the resuscitation of actively bleeding patients on warfarin.

Endoscopic therapy

A variety of endoscopic methods are available to treat PUB, with new devices coming to the market regularly. Guidelines recommend the use of injection therapy plus either thermal coagulation or endoclips [6]. Systematic reviews have confirmed that combined therapy is more effective than adrenaline injection alone, but it has not been shown that combined therapy is better than clips or thermal coagulation alone and these two modalities alone were found to be equivalent [7-9].

Two recent studies have examined the efficacy of the newly developed triclíp: in both a simulator model [10] and clinical practice, results were worse than with standard clips, mainly because of difficulty in placing this clip accurately [11]. A systematic review confirmed that endoclips are less effective for PUB lesions on the posterior duodenal wall, posterior gastric wall, and lesser curve because of difficulty placing the clips [8].

Adrenaline injection is probably the easiest and most widely available endoscopic technique and may be used alone when other methods are unavailable. The optimal volume of adrenaline for combination therapies is unclear, but recent studies have clarified the dose when used alone. Liou *et al.* [12] compared 20, 30 and 40 ml of 1:10,000 adrenaline (epinephrine). Initial haemostasis was equivalent but the smallest volume was associated with significantly more re-bleeding and the 40 ml dose was associated with an unacceptable perforation rate (6.6%), suggesting that when used alone, 30 ml would be the optimal dose. Injection regimes were further refined when 40 ml 1:10,000 adrenaline alone was compared with 40 ml saline alone or enough adrenaline to produce initial haemostasis followed by saline to a total volume of 40 ml. Initial haemostasis was higher in both the adrenaline groups (both 98.6%) than the saline alone group (88.9%) but re-bleeding rates were similar (approximately 5%). Perforation occurred in 2.8% of the adrenaline alone group [13]. These volumes are rather higher than most Western endoscopists generally use, but these data should encourage a re-examination of these practices. Although multimodality endoscopic therapies are preferable, when other methods are unavailable, adrenaline followed by saline would seem to be appropriate.

Pharmacotherapy

Potent acid suppressive therapy to stabilise the fibrin clot in the acute situation has always been an attractive idea and proton pump inhibitors (PPIs) are widely used for this. However, data on the most effective way to use these agents are conflicting. High dose intravenous omeprazole (80 mg bolus followed by 8 mg/h for 72 hours) after endoscopic therapy has been widely used following the

landmark study of Lau *et al.* [14]. However the effects of pharmacodynamics, physiology and *Helicobacter pylori* infection in the Asian population in that study mean that it may not be generalisable to other groups. A Cochrane systematic review (including the data from Lau *et al.*) confirmed that PPIs reduce re-bleeding and surgery in PUB but did not show the superiority of any dose or route of administration [15]. The beneficial effects were significantly more pronounced in the studies in Asian populations. One important point is that all the studies in the systematic review used oral or parenteral PPIs at higher than standard ulcer healing doses [15].

Subsequent recent studies examining the effects after endoscopic therapy have only added to the confusion. A retrospective Canadian study of 162 patients [16] showed that oral omeprazole (at least 40 mg) was as effective as omeprazole infusion following endoscopic haemostasis. A North American trial comparing parenteral high-dose pantoprazole with intravenous ranitidine was stopped early because of slow recruitment [17]; no significant differences were seen, although re-bleeding was numerically lower with pantoprazole (6.9 versus 14.3%). A similar randomized study predominantly in Europeans showed that intravenous pantoprazole was not superior to intravenous ranitidine, although re-bleeding rates in that study were very low (2.9-3.2%) [18]. A further trial in India (203 patients) showed that intravenous pantoprazole was superior to placebo, reducing re-bleeding (7.8 versus 19.8%) and transfusions [19]. A Greek study of 164 patients showed that intravenous pantoprazole was more effective than somatostatin in preventing re-bleeding (5 versus 17%) despite equivalent actions on gastric pH [20]. Thus, it seems that PPI therapy is mandatory for PUB and whilst high dose infusions are very effective in Asian populations, the optimal regimens for Caucasians still need to be established. A recently published large study [21] showed that intravenous esomeprazole bolus (80 mg) followed by infusion for 72 hours (8 mg/h) did significantly reduce re-bleeding (5.9 versus 10.3%) and endoscopic retreatment (6.4 versus 11.6%) compared to placebo in a predominantly Caucasian population (767 patients) and this may yet become the treatment of choice following endoscopic therapy.

The effect of potent acid suppression before endoscopy has also been investigated. Lau *et al.* [22] showed that parenteral omeprazole reduced endoscopic stigmata and need for intervention, but a further Cochrane analysis confirmed only that pre-endoscopy PPI reduced high-risk endoscopic stigmata without affecting the clinically important outcomes (mortality, re-bleeding or surgery) [23].

Prevention

It is accepted that *H. pylori*, cyclo-oxygenase inhibitors and antiplatelet drugs are the main remedial causes of PUB. Gisbert *et al.* [24] confirmed the importance of *H. pylori*. Following successful eradication (which may require sequential treatment strategies) the rate of re-bleeding was only 0.22% per year and patients did not require long-term acid suppressive therapy. Against this, a systematic review of *H. pylori* testing in PUB [25] showed that all tests except urea-breath testing (sensitivity 93%), especially biopsy-based ones (sensitivities 45-70%), had decreased sensitivity in PUB. Faecal antigen testing performed relatively well (sensitivity 87%). These studies confirm that strategies to identify, treat and follow up *H. pylori* are essential in the management of PUB.

Although the initial promise of the cyclo-oxygenase-2 (COX-2) selective agents has been tempered by worries about their cardiovascular toxicity and the realisation that they are not without gastrointestinal effects, these agents do offer some advantages in selected patients with regards to PUB. In the colonic adenoma prevention trial, rofecoxib (now withdrawn, but a very selective COX-2 inhibitor) was associated with PUB rates significantly greater than placebo (0.23 versus 0.06 per 100 patients-years), showing these are not a panacea for peptic ulcers [26]. However, a large population-based case-control study confirmed that COX-2 selective agents were associated with lower rates of PUB than traditional nonsteroidal anti-inflammatory drugs (NSAIDs) and residual risk was significantly reduced by co-prescription of a PPI [27].

Previously, celecoxib was shown to be equivalent to diclofenac plus PPI following PUB (re-bleeding rates 4.9 versus 6.4%) [28] and an important study, again from Hong Kong, showed that celecoxib plus esomeprazole was better than celecoxib alone following PUB (re-bleeding rates 0 versus 8.9%) [29]. There still may be differences between the various cyclo-oxygenase inhibitors but this regimen does offer a way forward for those requiring continued anti-inflammatory therapy after PUB.

Clopidogrel is associated with a significantly increased rate of PUB. Following an aspirin-induced PUB, clopidogrel alone was worse than aspirin plus PPI (re-bleeding rates after 52 weeks, 13.6 versus 0%) [30].

The addition of clopidogrel to aspirin does increase the absolute risk of PUB in those without a previous history of the condition, by about 1% per year (0.3% increased

to 1.4%) [31]. In view of this relatively low excess risk and the recent data suggesting that PPIs can significantly reduce the cardioprotective effects of clopidogrel [32], and the lack of proven efficacy for H₂-receptor antagonists in this situation [33], primary prophylaxis with a PPI is probably not indicated.

A difficult situation arises in those patients with PUB and a recently inserted coronary stent or unstable coronary syndrome. In these patients, discontinuation of dual antiplatelet agents is associated with a significant risk of adverse cardiovascular events (29% at 1 month [34]). In a small (88 patients) but important study, Ng *et al.* [35] showed that when antiplatelets were reintroduced within 1 week of PUB with PPI cover, the re-bleeding rate was 0% at 3 months. These results require duplication but suggest that antiplatelets can be reintroduced early after cessation of PUB.

Implications for clinical practice

The GBS score provides a secure evidence base to limit hospital admissions and investigations in low-risk patients. Combined endoscopic therapies are superior to adrenaline alone. When endoclips or thermocoagulation are unavailable, initial adrenaline followed by saline to a total volume of 40 ml seems the optimal injection regimen. Following cessation of bleeding, continued management should include searching for and eradicating *H. pylori* and strategies to minimise further drug-induced PUB. Aspirin combined with a PPI is preferable to clopidogrel following aspirin-induced PUB, and should anti-inflammatory drugs need to be reintroduced in this high-risk group, celecoxib plus esomeprazole seems to be the regimen of choice, bearing in mind cardiovascular risk factors. In the highest cardiovascular risk patients, early reintroduction of antiplatelets seems relatively safe, although further confirmatory studies are required as are further data on the interactions between antiplatelets and PPIs.

PPIs are mandatory after endoscopy in PUB, but the optimal regimen and drug do require further definition, especially in non-Asian patients. The best evidence is for high dose intravenous esomeprazole in this group of patients. High-dose PPI therapy before endoscopy does not affect major clinical outcomes, and should probably be reserved for patients in whom expedient endoscopy is not possible.

Abbreviations

COX-2, cyclo-oxygenase-2; GBS, Glasgow-Blatchford score; NSAID, nonsteroidal anti-inflammatory drug; PPI, proton pump inhibitor; PUB, peptic ulcer bleeding.

Competing interests

The author has received research funding from and served on advisory boards for AstraZeneca, and received funding for educational activities from AstraZeneca, Wyeth Pharmaceuticals and Abbott Pharmaceuticals, all of which manufacture PPIs.

References

- UK Comparative Audit of Upper Gastrointestinal Bleeding and the Use of Blood Project Group: *UK Comparative Audit of Upper Gastrointestinal Bleeding and the Use of Blood*. London, UK: British Society of Gastroenterology; 2007. [http://bsg.org.uk/pdf_word_docs/blood_audit_report_07.pdf].
 - Church NI, Dallal HJ, Masson J, Mowat NA, Johnston DA, Radin E, Turner M, Fullarton G, Prescott RJ, Palmer KR: **Validity of the Rockall scoring system after endoscopic therapy for bleeding peptic ulcer: a prospective cohort study.** *Gastrointest Endosc* 2006, **63**:606-12.
 - Stanley AJ, Ashley D, Dalton HR, Mowat C, Gaya DR, Thompson E, Warshaw U, Groome M, Cahill A, Benson G, Blatchford O, Murray W: **Outpatient management of patients with low-risk upper-gastrointestinal haemorrhage: multicentre validation and prospective evaluation.** *Lancet* 2009, **373**:42-7.
- Changes Clinical Practice**
 F1000 Factor 4.8 Must Read
 Evaluated by Ian Beales 26 Feb 2009, John Inadomi 03 Mar 2009
- Romagnuolo J, Barkun AN, Enns R, Armstrong D, Gregor J: **Simple clinical predictors may obviate urgent endoscopy in selected patients with nonvariceal upper gastrointestinal tract bleeding.** *Arch Intern Med* 2007, **167**:265-70.
 - Pabinger I, Brenner B, Kalina U, Knaub S, Nagy A, Ostermann H: **Prothrombin complex concentrate (Beriplex P/N) for emergency anticoagulation reversal: a prospective multinational clinical trial.** *J Thromb Haemost* 2008, **6**:622-31.
 - Scottish Intercollegiate Guidelines Network (SIGN): *Management of Acute Upper and Lower Gastrointestinal Bleeding. A National Clinical Guideline*. Edinburgh, UK: SIGN; Sept 2008. [<http://www.sign.ac.uk/pdf/sign105.pdf>].
 - Vergara M, Calvet X, Gisbert JP: **Epinephrine injection versus epinephrine injection and a second endoscopic method in high risk bleeding ulcers.** *Cochrane Database Syst Rev* 2007: CD005584.
 - Sung JJ, Tsoi KK, Lai LH, Wu JC, Lau JY: **Endoscopic clipping versus injection and thermo-coagulation in the treatment of non-variceal upper gastrointestinal bleeding: a meta-analysis.** *Gut* 2007, **56**:1364-73.
- Changes Clinical Practice**
 F1000 Factor 6.0 Must Read
 Evaluated by Ian Beales 09 Apr 2008
- Marmo R, Rotondano G, Piscopo R, Bianco MA, D'Angella R, Cipolletta L: **Dual therapy versus monotherapy in the endoscopic treatment of high-risk bleeding ulcers: a meta-analysis of controlled trials.** *Am J Gastroenterol* 2007, **102**:279-89.
 - Maiss J, Dumser C, Zopf Y, Naegel A, Krauss N, Hochberger J, Matthes K, Hahn EG, Schwab D: **"Hemodynamic efficacy" of two endoscopic clip devices used in the treatment of bleeding vessels, tested in an experimental setting using the compact Erlangen Active Simulator for Interventional Endoscopy (compactEASIE) training model.** *Endoscopy* 2006, **38**:575-80.
 - Lin HJ, Lo WC, Cheng YC, Perng CL: **Endoscopic hemoclip versus triclclip placement in patients with high-risk peptic ulcer bleeding.** *Am J Gastroenterol* 2007, **102**:539-43.
 - Liou TC, Lin SC, Wang HY, Chang WH: **Optimal injection volume of epinephrine for endoscopic treatment of peptic ulcer bleeding.** *World J Gastroenterol* 2006, **12**:3108-13.
 - Liou TC, Chang WH, Wang HY, Lin SC, Shih SC: **Large-volume endoscopic injection of epinephrine plus normal saline for peptic ulcer bleeding.** *J Gastroenterol Hepatol* 2007, **22**:996-1002.
 - Lau JY, Sung JJ, Lee KK, Yung MY, Wong SK, Wu JC, Chan FK, Ng EK, You JH, Lee CW, Chan AC, Chung SC: **Effect of intravenous omeprazole on recurrent bleeding after endoscopic treatment of bleeding peptic ulcers.** *N Engl J Med* 2000, **343**:310-6.
 - Leontiadis GI, Sharma VK, Howden CW: **Proton pump inhibitor therapy for peptic ulcer bleeding: Cochrane collaboration meta-analysis of randomized controlled trials.** *Mayo Clin Proc* 2007, **82**:286-96.
 - Murthy S, Keyvani L, Leeson S, Targownik LE: **Intravenous versus high-dose oral proton pump inhibitor therapy after endoscopic hemostasis of high-risk lesions in patients with acute nonvariceal upper gastrointestinal bleeding.** *Dig Dis Sci* 2007, **52**:1685-90.
 - Jensen DM, Pace SC, Soffer E, Comer GM; 315 Study Group: **Continuous infusion of pantoprazole versus ranitidine for prevention of ulcer rebleeding: a U.S. multicenter randomized, double-blind study.** *Am J Gastroenterol* 2006, **101**:1991-9; quiz 2170.
 - van Rensburg C, Barkun AN, Racz I, Fedorak R, Bornman PC, Beglinger C, Balanzo J, Deviere J, Kupcinskis L, Luehmann R, Doerfler H, Schafer-Preuss S: **Clinical trial: intravenous pantoprazole vs. ranitidine for the prevention of peptic ulcer re-bleeding: a multicentre, multinational, randomized trial.** *Aliment Pharmacol Ther* 2009, **29**:497-507.
 - Zargar SA, Javid G, Khan BA, Yattoo GN, Shah AH, Gulzar GM, Sodhi JS, Mujeeb SA, Khan MA, Shah NA, Shafi HM: **Pantoprazole infusion as adjuvant therapy to endoscopic treatment in patients with peptic ulcer bleeding: prospective randomized controlled trial.** *J Gastroenterol Hepatol* 2006, **21**:716-21.
 - Tsibouris P, Zintzaras E, Lappas C, Moussia M, Tsianos G, Galeas T, Potamianos S: **High-dose pantoprazole continuous infusion is superior to somatostatin after endoscopic hemostasis in patients with peptic ulcer bleeding.** *Am J Gastroenterol* 2007, **102**:1192-9.
 - Sung JJ, Barkun A, Kuipers EJ, Mossner J, Jensen D, Stuart R, Lau J, Ahlborn H, Kilhann J, Lind T: **Intravenous esomeprazole for prevention of recurrent peptic ulcer bleeding.** *Ann Int Med* 2009, **150**:455-464.
- Changes Clinical Practice**
 F1000 Factor 4.8 Must Read
 Evaluated by Ian Beales 06 May 2009, Philip O Katz 06 May 2009
- Lau JY, Leung WK, Wu JC, Chan FK, Wong VW, Chiu PW, Lee VW, Lee KK, Cheung FK, Siu P, Ng EK, Sung JJ: **Omeprazole before endoscopy in patients with gastrointestinal bleeding.** *N Engl J Med* 2007, **356**:1631-40.
- F1000 Factor 6.4 Must Read**
 Evaluated by Ian Beales 16 May 2007, Pankaj Jay Pasricha 21 May 2007
- Dorward S, Sreedharan A, Leontiadis GI, Howden CW, Moayyedi P, Forman D: **Proton pump inhibitor treatment initiated prior to endoscopic diagnosis in upper gastrointestinal bleeding.** *Cochrane Database Syst Rev* 2006:CD005415.
 - Gisbert JP, Calvet X, Feu F, Bory F, Cosme A, Almela P, Santolaria S, Aznarez R, Castro M, Fernandez N, Garcia-Gravalos R, Canete N, Benages A, Montoro M, Borda F, Perez-Aisa A, Pique JM: **Eradication of *Helicobacter pylori* for the prevention of peptic ulcer re-bleeding.** *Helicobacter* 2007, **12**:279-86.
 - Gisbert JP, Abaira V: **Accuracy of *Helicobacter pylori* diagnostic tests in patients with bleeding peptic ulcer: a systematic review and meta-analysis.** *Am J Gastroenterol* 2006, **101**:848-63.
 - Lanas A, Baron JA, Sandler RS, Horgan K, Bolognese J, Oxenius B, Quan H, Watson D, Cook TJ, Schoen R, Burke C, Loftus S, Niv Y, Ridell R, Morton D, Bresalier R: **Peptic ulcer and bleeding events associated with rofecoxib in a 3-year colorectal adenoma chemoprevention trial.** *Gastroenterology* 2007, **132**:490-7.

27. Garcia Rodriguez LA, Barreales Tolosa L: **Risk of upper gastrointestinal complications among users of traditional NSAIDs and COXIBs in the general population.** *Gastroenterology* 2007, **132**:498-506.

F1000 Factor 4.8 *Must Read*

Evaluated by Ian Beales 22 Mar 2007, Peter Katelaris 11 Apr 2007

28. Chan FK, Hung LC, Suen BY, Wu JC, Lee KC, Leung VK, Hui AJ, To KF, Leung WK, Wong VW, Chung SC, Sung JJ: **Celecoxib versus diclofenac and omeprazole in reducing the risk of recurrent ulcer bleeding in patients with arthritis.** *N Engl J Med* 2002, **347**:2104-10.
29. Chan FK, Wong VW, Suen BY, Wu JC, Ching JY, Hung LC, Hui AJ, Leung VK, Lee VW, Lai LH, Wong GL, Chow DK, To KF, Leung WK, Chiu PW, Lee YT, Lau JY, Chan HL, Ng EK, Sung JJ: **Combination of a cyclo-oxygenase-2 inhibitor and a proton-pump inhibitor for prevention of recurrent ulcer bleeding in patients at very high risk: a double-blind, randomised trial.** *Lancet* 2007, **369**:1621-6.
30. Lai KC, Chu KM, Hui WM, Wong BC, Hung WK, Loo CK, Hu WH, Chan AO, Kwok KF, Fung TT, Wong J, Lam SK: **Esomeprazole with aspirin versus clopidogrel for prevention of recurrent gastrointestinal ulcer complications.** *Clin Gastroenterol Hepatol* 2006, **4**:860-5.
31. Aronow HD, Steinhubl SR, Brennan DM, Berger PB, Topol EJ, CREDO Investigators: **Bleeding risk associated with 1 year of dual antiplatelet therapy after percutaneous coronary**

intervention: Insights from the Clopidogrel for the Reduction of Events During Observation (CREDO) trial. *Am Heart J* 2009, **157**:369-74.

Changes Clinical Practice

F1000 Factor 6.0 *Must Read*

Evaluated by Ian Beales 03 Apr 2009

32. Ho PM, Maddox TM, Wang L, Fihn S, Jesse R, Peterson E, Rumsfeld J: **Risk of adverse outcomes associated with concomitant use of clopidogrel and proton pump inhibitors following acute coronary syndrome.** *JAMA* 2009, **301**:937-44.

Changes Clinical Practice

F1000 Factor 8.1 *Exceptional*

Evaluated by Gregory Lip 12 Mar 2009, Marrick Kucin 23 Mar 2009, Melvin Cheitlin 08 Apr 2009

33. Ibáñez L, Vidal X, Vendrell L, Moretti U, Laporte JR: **Upper gastrointestinal bleeding associated with antiplatelet drugs.** *Aliment Pharmacol Ther* 2006, **23**:235-42.
34. Henriksen PA, Palmer K, Boon NA: **Management of upper gastrointestinal haemorrhage complicating dual anti-platelet therapy.** *QJM* 2008, **101**:261-7.
35. Ng FH, Chan P, Kwanching CP, Loo CK, Cheung TK, Wong SY, Kng C, Ng KM, Lai ST, Wong BC: **Management and outcome of peptic ulcers or erosions in patients receiving a combination of aspirin plus clopidogrel.** *J Gastroenterol* 2008, **43**:679-86.