Recent advances in pharmacological treatment of neuropathic pain

Nanna Brix Finnerup^{1*}, Søren Hein Sindrup² and Troels Staehelin Jensen¹

Addresses: ¹Danish Pain Research Center, Aarhus University Hospital, Norrebrogade 44, 8000 Aarhus C, Denmark; ²Department of Neurology, Odense University Hospital, Sdr. Boulevard 29, 5000 Odense, Denmark

* Corresponding author: Nanna Brix Finnerup (finnerup@ki.au.dk)

FI000 Medicine Reports 2010, **2**:52 (doi:10.3410/M2-52)

The electronic version of this article is the complete one and can be found at: <http://fi000.com/reports/medicine/content/2/52>

Abstract

Recent studies investigating the pharmacological management of neuropathic pain support the efficacy of certain antidepressants, anticonvulsants, and opioids. Novel directions in drug applications include topical applications of patches with either lidocaine or capsaicin and intradermal injections of botulinum toxin. In cases of partial pain relief, drug combinations may be considered.

Introduction and context

Neuropathic pain is a heterogeneous group of chronic pain conditions caused by lesion or disease of the peripheral or central nervous system [1]. Common neuropathic pain syndromes include painful polyneuropathy (e.g., due to diabetes), postherpetic neuralgia, peripheral nerve injury, radiculopathy, cancer-related pain, and central post-stroke pain. Neuropathic pain often has a negative impact on the mental health and quality of life in these groups of patients. Recommended first-line treatments are the anticonvulsants gabapentin and pregabalin, certain antidepressants (tricyclic antidepressants [TCAs] or serotonin-noradrenaline reuptake inhibitors [SNRIs]), and topical lidocaine [2,3]. Patients may, however, experience no or only partial pain reduction in tolerated doses.

Recent advances

Anticonvulsants and antidepressants

Different types of neuropathic pain may respond differently to a given drug and there is a need for studies where predictive factors for response are identified. In a recent well-designed randomized controlled trial (RCT), gabapentin in doses up to 2400 mg/day failed to relieve pain in patients with post-traumatic or post-operative nerve injury [4], although the study showed marked improvement on several secondary outcome measures, including pain relief and Patient Global Impression of Change. Also, flexible-dose regimes of pregabalin seem

to provide a better combination of pain relief and few discontinuations than a fixed dose regime [5].

Levetiracetam is a novel anti-epileptic drug that binds to the synaptic vesicle protein SV2A in the brain and spinal cord. Despite the antinociceptive effect found in animal models of neuropathic pain, levetiracetam did not relieve post-mastectomy syndrome and spinal cord injury pain [6,7].

The need for mechanism-based approaches and identification of possible predictors for response is also relevant for sodium channel blockers. Except for the use of sodium channel blockers in trigeminal neuralgia, where they are first-line medication, there is limited evidence for efficacy of drugs like oxcarbazepine, lamotrigine, and the newer anticonvulsant lacosamide [2,8,9] in neuropathic pain. However, despite the lack of an overall effect in large-scale trials, subgroups of patients do seem to respond.

Studies with head-to-head comparisons find only minor differences in efficacy between TCAs and gabapentin/pregabalin [10]. SNRIs are also effective in patients with painful polyneuropathy [11] and can be an alternative to TCAs, in particular in patients with cardiac disorders. Selective serotonin reuptake inhibitors provide a clinically meaningful effect in only few patients [12].

Opioids

Opioids also have a role in the treatment of neuropathic pain. Their efficacy is comparable to that of gabapentin/pregabalin and TCAs [2,13,14], but they are not considered first-line drugs due to side effects, tolerance, uncertain long-term safety and efficacy, and so on.

Combination therapy

In cases of partial but insufficient pain relief by a single drug, combinations are often used. Due to a lack of controlled studies, the rationale for such combination therapy has mainly been based on theory, but is now supported by recent RCTs. The best evidence is for the combination of a TCA or an opioid with gabapentin [15-17]. Such combinations are suggested to improve treatment compared with treatment with each drug alone in maximum tolerated doses.

New drug classes

Topical lidocaine is a safe treatment with no or limited systemic side effects. New evidence from an open-label study suggests that lidocaine patches are useful, not only in postherpetic neuralgia or very localized neuropathic pain but also in painful diabetic neuropathy [18].

Transient receptor TRPV1 agonists and antagonists represent a new class of analgesics [19]. A single application of a high-concentration capsaicin dermal patch produced sustained pain reduction in patients with postherpetic neuralgia [20] and painful HIV-associated sensory polyneuropathy [21], although the long-term effect and the frequency of treatments have not been examined. NGX-4010 (marketed name Qutenza®), a cutaneous patch of capsaicin 8%, has been given marketing authorization in Europe. The approved indication is: "treatment of peripheral neuropathic pain in non-diabetic adults". NGX-4010 treatments may be repeated every 90 days.

Botulinum toxin type A (BTX-A) given intradermally is another novel topical treatment approach that has been shown to relieve focal painful neuropathy [22], and painful diabetic polyneuropathy in two RCTs [23]. Patients received multiple intradermal injections once in the affected area and the pain reduction lasted for 12 weeks. Large-scale trials and examination of long-term efficacy are needed to test its value for clinical practice.

A phase 2 RCT found that the neuronal nicotinic acetylcholine receptor agonist ABT-594 significantly reduced pain intensity compared with placebo in patients with painful diabetic neuropathy [24]. Unfortunately, ABT-594 treatment was associated with a high number of intolerable dose-related side effects, but the

study suggests that further development of this drug class may present new therapeutic options.

Implications for clinical practice

The best available evidence and knowledge of side-effect profiles still support the current guidelines of using TCAs (or alternatively SNRIs), gabapentin, and pregabalin as first-line therapy and opioids as second line therapy for neuropathic pain. The side-effect profile for opioids is judged differently by some researchers, which may explain why opioids are considered by some as first-line therapy for certain conditions. In order to minimize side effects, recent advances have led to the development of topical agents, including lidocaine patches, NGX-4010 (high-dose capsaicin), and intradermal BTX-A. There is not abundant evidence for these treatments, but their advantage is that they have no or only minor systemic side effects.

If one drug provides only partial pain relief, a combination with another drug class is recommended – for example, an antidepressant or an opioid in combination with gabapentin or pregabalin, or a combination of a topical agent with an oral drug.

Carbamazepine, oxcarbazepine, lamotrigine, and lacosamide have no primary role in the treatment of neuropathic pain (except for trigeminal neuralgia), but whether we can identify subgroups of patients with specific symptoms or signs who will benefit from these treatments is still to be determined. This emphasizes the need for a mechanism-based approach, where patients are classified based on constellations of symptoms and signs rather than a disease-based classification. Hopefully, this will lead to better patient-oriented treatment. Levetiracetam does not seem to have a role in the treatment of neuropathic pain.

Abbreviations

BTX-A, botulinum toxin type A; RCT, randomized controlled trial; SNRI, serotonin noradrenaline reuptake inhibitor; TCA, tricyclic antidepressant.

Competing interests

Within the past two years, NBF has received research support and/or honoraria from UCB-Nordic and Grünenthal GmbH, TSJ has received research support and/or advisory fees from Pfizer, Grünenthal, Esteve, Glaxo-SmithKline, and Eli Lilly, and SHS has received research support from UCB-Nordic and advisory fees from Grünenthal and Eli Lilly.

Acknowledgements

The work behind this review was supported by the Velux Foundation. We would like to thank research secretary Helle Obenhausen Andersen for language revision.

References

1. Treede RD, Jensen TS, Campbell JN, Cruccu G, Dostrovsky JO, Griffin JW, Hansson P, Hughes R, Nurmikko T, Serra J: **Neuropathic pain: redefinition and a grading system for clinical and research purposes.** *Neurology* 2008, **70**:1630-5.
2. Finnerup NB, Otto M, McQuay HJ, Jensen TS, Sindrup SH: **Algorithm for neuropathic pain treatment: an evidence based proposal.** *Pain* 2005, **118**:289-305.

F1000 Factor 6.0 Must Read
Evaluated by Michael Serpell 14 Mar 2006
3. Dworkin RH, O'Connor AB, Backonja M, Farrar JT, Finnerup NB, Jensen TS, Kalso EA, Loeser JD, Miasowski C, Nurmikko TJ, Portenoy RK, Rice AS, Stacey BR, Treede RD, Turk DC, Wallace MS: **Pharmacologic management of neuropathic pain: Evidence-based recommendations.** *Pain* 2007, **132**:237-51.

F1000 Factor 6.0 Must Read
Evaluated by Andreas Kopf 16 Jul 2008
4. Gordh TE, Stubhaug A, Jensen TS, Arner S, Biber B, Boivie J, Mannheimer C, Kalliomaki J, Kalso E: **Gabapentin in traumatic nerve injury pain: A randomized, double-blind, placebo-controlled, cross-over, multi-center study.** *Pain* 2008, **138**:255-66.

F1000 Factor 3.0 Recommended
Evaluated by Claudia Sommer 09 Jan 2009
5. Stacey BR, Barrett JA, Whalen E, Phillips KF, Rowbotham MC: **Pregabalin for postherpetic neuralgia: placebo-controlled trial of fixed and flexible dosing regimens on allodynia and time to onset of pain relief.** *J Pain* 2008, **9**:1006-17.

F1000 Factor 3.0 Recommended
Evaluated by Kenneth Candido 12 Mar 2009
6. Vilholm OJ, Cold S, Rasmussen L, Sindrup SH: **Effect of levetiracetam on the postmastectomy pain syndrome.** *Eur J Neurol* 2008, **15**:851-7.

F1000 Factor 3.0 Recommended
Evaluated by Nanna Finnerup 14 Oct 2008
7. Finnerup NB, Grydehoj J, Bing J, Johannesen IL, Biering-Sorensen F, Sindrup SH, Jensen TS: **Levetiracetam in spinal cord injury pain: a randomized controlled trial.** *Spinal Cord* 2009, **47**:861-7.
8. Rao RD, Flynn PJ, Sloan JA, Wong GY, Novotny P, Johnson DB, Gross HM, Renno SI, Nashawaty M, Loprinzi CL: **Efficacy of lamotrigine in the management of chemotherapy-induced peripheral neuropathy: a phase 3 randomized, double-blind, placebo-controlled trial, N01C3.** *Cancer* 2008, **112**:2802-8.
9. Wymer JP, Simpson J, Sen D, Bongardt S, Lacosamide SP742 Study Group: **Efficacy and safety of lacosamide in diabetic neuropathic pain: an 18-week double-blind placebo-controlled trial of fixed-dose regimens.** *Clin J Pain* 2009, **25**:376-85.
10. Bansal D, Bhansali A, Hota D, Chakrabarti A, Dutta P: **Amitriptyline vs. pregabalin in painful diabetic neuropathy: a randomized double blind clinical trial.** *Diabet Med* 2009, **26**:1019-26.
11. Kajdasz DK, Iyengar S, Desai D, Backonja MM, Farrar JT, Fishbain DA, Jensen TS, Rowbotham MC, Sang CN, Ziegler D, McQuay HJ: **Duloxetine for the management of diabetic peripheral neuropathic pain: evidence-based findings from post hoc analysis of three multicenter, randomized, double-blind, placebo-controlled, parallel-group studies.** *Clin Ther* 2007, **29**(Suppl):2536-46.

F1000 Factor 3.0 Recommended
Evaluated by Stephen Tyrer 10 Mar 2008
12. Otto M, Bach FW, Jensen TS, Brosen K, Sindrup SH: **Escitalopram in painful polyneuropathy: A randomized, placebo-controlled, cross-over trial.** *Pain* 2008, **139**:275-83.
13. Wu CL, Agarwal S, Tella PK, Klick B, Clark MR, Haythornthwaite JA, Max MB, Raja SN: **Morphine versus mexiletine for treatment of postamputation pain: a randomized, placebo-controlled, crossover trial.** *Anesthesiology* 2008, **109**:289-96.

F1000 Factor 3.0 Recommended
Evaluated by James Eisenach 17 Sep 2008
14. Norrbrink C, Lundeborg T: **Tramadol in neuropathic pain after spinal cord injury: a randomized, double-blind, placebo-controlled trial.** *Clin J Pain* 2009, **25**:177-84.
15. Gilron I, Bailey JM, Tu D, Holden RR, Weaver DF, Houlden RL: **Morphine, gabapentin, or their combination for neuropathic pain.** *N Engl J Med* 2005, **352**:1324-34.

Changes Clinical Practice
F1000 Factor 6.5 Must Read
Evaluated by Howard Gutstein 16 Nov 2005, Michael Serpell 12 Dec 2005, Nadine Attal 13 Feb 2006
16. Hanna M, O'Brien C, Wilson MC: **Prolonged-release oxycodone enhances the effects of existing gabapentin therapy in painful diabetic neuropathy patients.** *Eur J Pain* 2008, **12**:804-13.

F1000 Factor 3.0 Recommended
Evaluated by Claudia Sommer 14 Aug 2008
17. Gilron I, Bailey JM, Tu D, Holden RR, Jackson AC, Houlden RL: **Nortriptyline and gabapentin, alone and in combination for neuropathic pain: a double-blind, randomised controlled crossover trial.** *Lancet* 2009, **374**:1252-61.

F1000 Factor 4.9 Must Read
Evaluated by Nanna Finnerup 02 Dec 2009, Joel Katz 11 Dec 2009, Didier Bouhassira 21 Dec 2009
18. Baron R, Mayoral V, Leijon G, Binder A, Steigerwald I, Serpell M: **5% lidocaine medicated plaster versus pregabalin in postherpetic neuralgia and diabetic polyneuropathy: an open-label, non-inferiority two-stage RCT study.** *Curr Med Res Opin* 2009, **25**:1663-76.
19. Wong GY, Gavva NR: **Therapeutic potential of vanilloid receptor TRPV1 agonists and antagonists as analgesics: Recent advances and setbacks.** *Brain Res Rev* 2009, **60**:267-77.

F1000 Factor 6.0 Must Read
Evaluated by Peter Gerner 06 May 2009
20. Backonja M, Wallace MS, Blonsky ER, Cutler BJ, Malan P, Jr., Rauck R, Tobias J: **NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study.** *Lancet Neurol* 2008, **7**:1106-12.
21. Simpson DM, Brown S, Tobias J: **Controlled trial of high-concentration capsaicin patch for treatment of painful HIV neuropathy.** *Neurology* 2008, **70**:2305-13.

F1000 Factor 3.0 Recommended
Evaluated by Thien Nguyen 24 Jun 2008
22. Ranoux D, Attal N, Morain F, Bouhassira D: **Botulinum toxin type A induces direct analgesic effects in chronic neuropathic pain.** *Ann Neurol* 2008, **64**:274-83.

F1000 Factor 4.8 Must Read
Evaluated by Troels Jensen 06 Jan 2009, Claudia Sommer 09 Jan 2009
23. Yuan RY, Sheu JJ, Yu JM, Chen WT, Tseng IJ, Chang HH, Hu CJ: **Botulinum toxin for diabetic neuropathic pain: a randomized double-blind crossover trial.** *Neurology* 2009, **72**:1473-8.

F1000 Factor 3.0 Recommended
Evaluated by A Gordon Smith 05 Jun 2009
24. Rowbotham MC, Rachel DW, Thomas J, Nothaft W, Backonja MM: **A randomized, double-blind, placebo-controlled trial evaluating the efficacy and safety of ABT-594 in patients with diabetic peripheral neuropathic pain.** *Pain* 2009, **146**:245-52.

F1000 Factor 3.0 Recommended
Evaluated by Nanna Finnerup 05 Oct 2009